

IL 12 Human

Description: Interleukin-12 Human Recombinant produced in HEK cells is a glycosylated heterodimer, having a total molecular weight of 57kDa. The IL12 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-108

For research use only.

Synonyms: NKSF, CTL maturation factor (TCMF), Cytotoxic lymphocyte maturation factor (CLMF), TSF, Edodekin-alpha, IL-12.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: IL-12 is a heterodimer of IL-12A and IL-12B linked through a disulfide-bond between cysteines in red in sequences below. >IL-12
ARNLPVATPDGPMFPCLLHHSQNLLRAVSNNMLQKARQTLEFYPTCTSEEIDHEDTKDKTSTVEACL
PLELTKNESCLNSRETSFITNGSCLSRKTSFMMALCLSSIEDLKMYQVEFKTMNAKL

Purity: Greater than 95% as observed by SDS-PAGE.

Formulation:

The IL12 was lyophilized from 1mg/ml in 1xPBS.

Stability:

Lyophilized IL-12 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL12 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL12 in sterile PBS containing 0.1% endotoxin-free recombinant HSA.

Introduction:

IL-12 is a heterodimeric cytokine that stimulates the production of interferon gamma from T-cells and natural killer cells, and also induces differentiation of Th1 helper cells. IL-12 is an initiator of cell-mediated immunity.

Biological Activity:

The specific activity was determined by the dose-dependent release of IFN-gamma from the human NK92 cell line in presence of 20ng/mL rIL-2, and is typically 0.1-0.5ng/ml.

To place an order, please [Click HERE](#).