

IL 12 Mouse

Description: Interleukin-12 Mouse Recombinant produced in Sf9 insect cells is a glycosylated disulfide linked heterodimeric polypeptide containing 506 amino acids and having a molecular weight of 75 kDa comprised of disulfide-bonded 35 kDa (p35) and 40 kDa (p40) subunits. The IL-12 is purified by proprietary chromatographic techniques.

Catalog #: CYP5-151

For research use only.

Synonyms: NKSF, CTL maturation factor (TCMF), Cytotoxic lymphocyte maturation factor (CLMF), TSF, Edodekin-alpha, IL-12.

Source: HEK-293 Cells.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: p35 Subunit: RVIPVSGPAR CLSQSRNLLK TTDDMVKTAR
EKLEKHYSCA EDIDHEDTR DQTSTLKTCL PLELHKNECS LATRETSSTT RGSCLPQKT
SLMMTLCLGS IYEDLKMYQT EFQAINAALQ NHNHQQIILD KGMLVAIDEL MQSLNHNGET
LRQKPPVGEA DPYRVKMKLC ILLHAFSTRV VTINRVMGYL SSA. p40 Subunit: MWELEKDVYV
VEVDW

Purity: Greater than 95% as determined by SDS-PAGE.

Formulation:

The protein was lyophilized from 0.5xPBS, pH 7.5.

Stability:

Lyophilized mouse IL-12 although stable at room temperature for 3 weeks, should be stored below -18°C. Upon reconstitution IL12 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized murine IL12 in sterile distilled pyrogen free water at 100

Introduction:

IL-12 is a heterodimeric cytokine that stimulates the production of interferon gamma from T-cells and natural killer cells, and also induces differentiation of Th1 helper cells. It is an initiator of cell-mediated immunity.

Biological Activity:

IL-12 Mouse has full biological activity when compared to standards. The ED50 is determined by the dose-dependent induced expression of CD25 on CD8+ mouse T cells is 10-20 ng/ml corresponding to a specific activity of 50,000-100,000 units/mg.

To place an order, please [Click HERE](#).