

4 1BBL Human

Description: 4 1BBL Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 184 amino acids and having a molecular mass of 19.5kDa. The 4 1BBL is purified by proprietary chromatographic techniques.

Synonyms: CD137L, CD137-L, 4-1BBL, 4-1BB Ligand, TNFSF9, Tumor Necrosis Factor (ligand) Superfamily Member 9.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: REGPELSPDD PAGLLDLRQG MFAQLVAQNV LLIDGPLSWY
SDPGLAGVSL TGGLSYKEDT KELVVAAGV YYVFFQLELR RVVAGEGSGS VSLALHLQPL
RSAAGAAALA LTVDLPPASS EARNSAFGFQ GRLLHLSAGQ RLGVLHTEA RARHAWQLTQ
GATVLGLFRV TPEIPAGLPS PRSE

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

Lyophilized from a 0.2m filtered concentrated solution in PBS, pH 7.4.

Stability:

Lyophilized 4 1BBL although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution 4 1BBL should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized 4 1BBL in sterile 18M-cm H₂O not less than 100

Introduction:

4-1BBL is a transmembrane cytokine that is part of the tumor necrosis factor (TNF) ligand family. 4-1BBL is a bidirectional signal transducer that performs as a ligand for TNFRSF9, which is a costimulatory receptor molecule in T lymphocytes. TNFSF9 and its TNFRSF9 take part in the antigen presentation development and in the generation of cytotoxic T cells. 4-1BBL is absent from resting T lymphocytes but rapidly expressed upon antigenic stimulation. TNFSF9 reactivates anergic T lymphocytes as well as promoting T lymphocyte proliferation. 4-1BB Ligand is needed for the optimal CD8 responses in CD8 T cells. 4-1BBL is expressed in carcinoma cell lines, and is thought to be involved in T cell-tumor cell interaction. 4-1BBL is expressed by activated B cells, macrophages, dendritic cells, activated T cells, neurons and astrocytes. The interaction of 4-1BB with TNFRSF9 strongly regulates immunity and has been proposed to preferentially control T cell responses based on studies in various murin.

Biological Activity:

Fully biologically active when compared to standard. Determined by the dose-dependent

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
stimulation of IL-8 production by human PBMC. The ED50 for this effect is 5-10ng/ml
corresponding to a Specific Activity of 100,000-200,000IU/mg.



To place an order, please [Click HERE](#).

Catalog #:CYPs-156

For research use only.

© 2015 NeoScientific

