

## MIF Human His C

**Description:** MIF human Recombinant, fused to His-tag at C-terminus, was cloned into an E. coli expression vector and was purified to apparent homogeneity by using conventional column chromatography techniques. Macrophage Inducing Factor Human Recombinant is a single, non-glycosylated, polypeptide chain containing 123 amino acids and having a molecular mass of 13.5 kDa.

**Catalog #:** CYP5-528

For research use only.

**Synonyms:** Phenylpyruvate tautomerase, Glycosylation-inhibiting factor, GIF, MMIF, MIF.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered lyophilized powder.

**Amino Acid Sequence:**

MPMFIVNTNVPASVDPDGLSELTTQLAQATGKPPQYIAVHVVPDQLMAFGGSSEPCALCSLHSI  
GKIGGAQNRSYSKLLCGLLAERLRISPDRVYINYYDMNAANVGWNNSTFALEHHHHHH.

**Purity:** Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

Human MIF was lyophilized from a 1mg/ml solution containing PBS pH-7.4.

**Stability:**

Lyophilized MIF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MIF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized MIF in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

The cytokine Macrophage Migration Inhibitor (MIF) has been identified to be secreted by the pituitary gland and the monocyte/macrophage and to play an important role in endotoxic shock. MIF has the unique property of being released from macrophages and T cells in response to physiological concentrations of glucocorticoids. The secretion of MIF is tightly regulated and decreases at high, anti-inflammatory steroid concentration.

**Biological Activity:**

Measured by its ability to bind rhCD74 in a functional ELISA.

**To place an order, please [Click HERE](#).**