www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

# HGF A Human

Description: HGF-A Human Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 463 amino acids fragment (32-494) having a total molecular weight of 57.8kDa. The HGF-A is fused with a 4.5kDa amino-terminal hexahistidine tag.The HGF-A is purified by proprietary chromatographic techniques.

Catalog #:CYPS-671

For research use only.

Synonyms: Scatter Factor, SF, Hepatopoietin, HPTA, HGF, HGFB, F-TCF, DFNB39, Hepatocyte growth factor, Hepatopoeitin-A, Hepatocyte growth factor alpha chain.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

#### Amino Acid Sequence:

QRKRRNTIHEFKKSAKTTLIKIDPALKIKTKKVNTADQCANRCTRNKGLPFTCKAFVFDKARKQCL WFPFNSMSSGVKKEFGHEFDLYENKDYIRNCIIGKGRSYKGTVSITKSGIKCQPWSSMIPHEHSFL PSSYRGKDLQENYCRNPRGEEGGPWCFTSNPEVRYEVCDIPQCSEVECMTCNGESYRGLMDH TESGKICORWDHOTPHRHKFI PERYPDKGEDDNYCRNPDGOPRPWCYTI DPHTRWEYCAIK

Purity: Greater than 95.0% as determined by SDS-PAGE.

## Formulation:

HGF-A protein is supplied in 22mM Na.Acetate pH4.8, 1mM EDTA and 50% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis. HGF is secreted as a single inactive polypeptide which is cleaved by serine proteases into a 69kDa Alpha chain and 34kDa Beta chain. A disulfide bond linking the alpha and beta chains produces the active, heterodimeric molecule.

To place an order, please Click HERE.





