

UCHL5 Human

Description: UCHL5 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 349 amino acids (1-329 a.a.) and having a molecular mass of 39.7 kDa. UCHL5 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: ENPS-104

For research use only.

Synonyms: Ubiquitin Carboxyl-terminal Hydrolase L5, UCH37, CGI-70, UCH-L5, INO80R, Ubiquitin thioesterase L5, INO80 complex subunit R, EC 3.4.19.12, Ubiquitin Carboxyl-terminal Esterase L5.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHSSGLVPRGSH MTGNAGEWCL MESDPGVFTE
LIKGFGRGA QVEEWSLEP ENFEKLKPVH GLIFLKWQP GEEPAGSVVQ DSRLDTIFFA
KQVINNACAT QAIVSVLLNC THQDVHLGET LSEFKEFSQS FDAAMKGLAL SNSDVIRQVH
NSFARQQMFE FDTKTSAKEE DAFHFVSYP VNGRLYELDG LREGPIDLGA CNQDDWISAV
RPVIEKRIQK YSE

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

The UCHL5 protein solution (0.5 mg/ml) is formulated in 20 mM Tris-HCl buffer (pH 8.0), 5 mM DTT, 200 mM NaCl, 0.1 mM PMSF, 2 mM EDTA and 30% glycerol.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

UCHL5 is a member of the peptidase C12 family. UCHL5 protein is a protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains, and deubiquitinating enzyme related to the 19S regulatory subunit of the 26S proteasome.

Biological Activity:

Specific activity: > 500 pmole/min/μg. Measured by the hydrolysis of Ubiquitin-AMC at pH 8.0, at 37°C.

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

To place an order, please [Click HERE](#).