

NTHL1 Human

Description: NTHL1 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 332 amino acids (1-312 a.a.) and having a molecular mass of 36.6kDa. NTHL1 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: ENPS-179

For research use only.

Synonyms: Endonuclease III-like protein 1, NTHL1, NTH1, OCTS3.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MCSPQESGMT ALSARMLTRS
RSLGPGAGPR GCREEPGLR RREAAAEARK SHSPVKRPRK AQRLRVAYEG SDSEKGEAE
PLKVPVWEPQ DWQQQLVNIR AMRNKKDAPV DHLGTEHCYD SSAPPKVRRY QVLLSLMLSS
QTKDQVTAGA MQLRLRAGLT VDSILQTDDA TLGKLIYPVG FWRSKVKYIK QTSAILQQHY
GGDIPASVAE LV

Purity: Greater than 80.0% as determined by SDS-PAGE.

Formulation:

NTHL1 protein solution (0.25mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 40% glycerol, 0.15M NaCl and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Nth endonuclease III-like protein 1 (NTHL1) is one of the enzymes which act on a large number of oxidatively damaged DNA sites. NTHL1 has an apurinic and/or apyrimidinic endonuclease activity and a DNA N-glycosylase activity. NTHL1 acts on a damaged strand, 5' from the damaged site and incises damaged DNA at cytosines, thymines and guanines.

To place an order, please [Click HERE](#).