www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

DsbC

Description: Disulfide-Bond Isomerase Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 216 amino acids (21-236) and having a molecular mass of 23.5 kDa.DsbC is purified by proprietary chromatographic techniques.

Catalog #:ENPS-298

For research use only.

Synonyms:DsbC, Thiol:disulfide interchange protein dsbC.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: DDAAIQQTLA KMGIKSSDIQ PAPVAGMKTV LTNSGVLYIT DDGKHIIQGP MYDVSGTAPVNVTNKMLLKQ LNALEKEMIV YKAPQEKHVI VFTDITCGY CHKLHEQMAD YNALGITVRYLAFPROGLDS DAEKEMKAIW CAKDKNKAFD DVMAGKSVAP ASCDVDIADH YVLGVQLGVSGTPAVVLSNG TLVPGYQPPK EMKEFLDEHQ KMTSGK.

Purity: Greater than 95.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

1 mg/ml solution containing 20mM phosphate buffer pH 7.5 and 20mM EDTA.

Stability:

Disulfide-Bond Isomerase Recombinant although stable at 4°C for 1 week, should be stored desiccated below -18°C.Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Dsb proteins (DsbA, DsbB, DsbC, and DsbD) catalyze formation and isomerization of protein disulfide bonds in the periplasm of Escherichia coli. DsbC is periplasmic enzyme known as a disulfide isomerase and can convert aberrant disulfide bonds to correct ones.

To place an order, please Click HERE.





