

ErbB2 Human

Description: ErbB-2 Human Recombinant is a 43.4 kDa protein containing 397 amino acid residues of the human Herstatin, and an extra Methionine at N-Terminal (underlined), produced in E.coli. The ErbB2 is purified by proprietary chromatographic techniques.

Synonyms: NEU, NGL, HER2, TKR1, HER-2, c-erb B2, HER-2/neu, Receptor tyrosine-protein kinase erbB-2, EC 2.7.10.1, p185erbB2, C-erbB-2, NEU proto-oncogene, Tyrosine kinase-type cell surface receptor HER2, MLN 19, CD340 antigen.

Source: Escherichia Coli.

Physical Appearance: Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MTQVCTGTDM KLRLPASPET HLDMLRHLYQ GCQVVQGNLE
LTYLPTNASL SFLQDIQEVQ GYVLIHNVQ RQVPLQLRLI VRGTQLFEDN YALAVLDNGD
PLNNTTPVTG ASPGGLRELQ LRSLTEILKG GVLIQRNPQL CYQDTILWKD IFHKNNQLAL
TLIDTNSRA CHPCSPMCKG SRCWGESSED CQSLTRTVCA GGCARCKGPL PTDCCHEQCA
AGCTGPKHSD CL

Purity: Greater than 95% as determined by SDS-PAGE.

Formulation:

Filtered (0.4

Stability:

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

ELISA, Western blotting.

Solubility:

It is recommended to add 0.1M Acetate buffer pH-4 to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10g/ml. In higher concentrations the solubility of this antigen is limited. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture. It is recommended to add 5mM DTT and 0.1-0.15M NaCl before freezing in order to prevent potential aggregation.

Introduction:

HER-2/neu (erbB-2) encodes an 185-kDa orphan receptor tyrosine kinase that is constitutively active as a dimer and displays potent oncogenic activity when overexpressed. Herstatin, as the product of alternative HER-2 transcript, retains intron 8. The herstatin mRNA is expressed in normal human fetal kidney and liver, but is at reduced levels relative to p185HER-2 mRNA in carcinoma cells that contain an amplified HER-2 gene. Herstatin appears to be an inhibitor of

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

p185HER-2, because it disrupts dimers, reduces tyrosine phosphorylation of p185, and inhibits the anchorage-independent growth of transformed cells that overexpress HER-2.



To place an order, please [Click HERE](#).

Catalog #:PKPS-350

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a
CATALOG
[CLICK HERE](#)