

S100A12 Human

Description: The Recombinant Human S100A12 produced in E.coli has a molecular mass of 11.63kDa containing 101 amino acid residues of the human S100A12 and fused to a 10 a.a. His tag at N-terminus.

Catalog #: PRPS-159

For research use only.

Synonyms: Extracellular newly identified RAGE-binding protein, Protein S100-A12, CGRP, Calcium-binding protein in amniotic fluid 1, CAAF1, Calgranulin-C, CAGC, EN-RAGE, Neutrophil S100 protein, S100 calcium-binding protein A12, p6, S100A12, MRP6, ENRAGE.

Source: Escherichia Coli.

Amino Acid Sequence: MKHHHHHHAS TKLEEHLEGI VNIFHQYSVR KGHFDTLSKG
ELKQLLTKEI ANTIKNIKDK AVIDEIFQGL DANQDEQVDFQEFISLVAIA LKAAHYHCHK E.

Formulation:

S100A12 was filtered (0.4

Stability:

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to add deionized water to prepare a working stock solution of approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

Introduction:

S100A12 belongs to the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 family members are localized in the cytoplasm and/or nucleus of a wide range of cells, and are involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100A12 may be involved in specific calcium-dependent signal transduction pathways and its regulatory effect on cytoskeletal components may modulate various neutrophil activities.

To place an order, please [Click HERE](#).