

PCNA Human

Description: PCNA Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 261 amino acids and having a molecular mass of 28769 Dalton.

Catalog #: PRPS-622

Synonyms: Proliferating cell nuclear antigen, PCNA, Cyclin, MGC8367.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MFEARLVQGS ILKKVLEALK DLINEACWDI SSSGVNLQSM
DSSHVSLVQLTLRSEGFDTY RCDRNLMAGV NLTSMSKILK CAGNEDIITL RAEDNADTLA
LVFEAPNQE K VSDYEMKLMD LDVEQLGIPE QEYSCVVKMP SGEFARICRD LSHIGDAVVI
SCAKDGVKFS ASGELGNGNI KLSQTSNVDK EEEAVTIEMN EPVQLTFALR YLNFFTKATP
LSSTVTLSMS ADV

Purity: Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The PCNA solution contains 20mM Tris pH-7.5 & 2mM EDTA & 20% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

PCNA is located in the nucleus and is a cofactor of DNA polymerase delta. PCNA acts as a homotrimer and helps elevate the processivity of leading strand synthesis during DNA replication. In reaction to DNA damage, PCNA is ubiquitinated and takes part in the RAD6-dependent DNA repair pathway. 2 transcript variants encoding the same protein have been found for PCNA gene. Pseudogenes of this PCNA gene have been described on chromosome 4 and on the X chromosome. PCNA is expressed during late G1- phase, S-phase of mitosis and persists until the end of the M-phase because of its long biological half-life. PCNA is induced by UV irradiation.

To place an order, please [Click HERE](#).