

ARF3 Human

Description: ARF3 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 201 amino acids (1-181 a.a.) and having a molecular mass of 22.8kDa. ARF3 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-940

For research use only.

Synonyms: ADP-ribosylation factor 3, ARF3.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGNIFGNLLK SLIGKKEMRI
LMVGLDAAGK TTILYKCLKG EIVTTIPTIG FNVETVEYKN ISFTVWDVGG QDKIRPLWRH
YFQNTQGLIF VVDSNDRERV NEAREELMRM LAEDELDAV LLVFANKQDL PNAMNAAEIT
DKLGLHSLRH RNWYIQATCA TSGDGLYEGL DWLANQLKNK K.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

ARF3 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 20% glycerol and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

ADP-ribosylation factor 3 (ARF3) belongs to the human ARF gene family. This family encodes small guanine nucleotide-binding proteins which stimulate the ADP-ribosyltransferase activity of cholera toxin and have a role in vesicular trafficking and as activators of phospholipase D. ARF3 functions as an allosteric activator of the cholera toxin catalytic subunit, an ADP-ribosyltransferase. ARF3 is involved in protein trafficking; may modulate vesicle budding and uncoating within the Golgi apparatus. The ARF3 gene is comprised of 5 exons and 4 introns.

To place an order, please [Click HERE](#).